

Four non-yellow species of *Rhizocarpon* new to China

LING HU¹*, XIAO ZHANG²*, CHUN-XIAO WANG², ZUN-TIAN ZHAO^{2*}

¹ Institute of Environment and Ecology, Shandong Normal University,
Jinan, 250014, P. R. China

² Key Laboratory of Plant Stress Research, College of Life Sciences,
Shandong Normal University, Jinan, 250014, P. R. China

*CORRESPONDENCE TO: ztzhao@sdu.edu.cn

ABSTRACT—Four species of the lichen genus *Rhizocarpon* (*R. cinereonigrum*, *R. distinctum*, *R. sublavatum*, and *R. subpostumum*) are reported for the first time from China. Descriptions provide morphological and chemical characters and known distribution of each species.

KEY WORDS—lichen-forming fungi, *Rhizocarpaceae*, saxicolous, taxonomy

Introduction

Rhizocarpon Ramond ex DC. (*Rhizocarpaceae*, *Rhizocarpales*, *Lecanoromycetes*, *Ascomycota*) was established by de Candolle in 1805 (Ihlen 2004). *Rhizocarpon* can best be characterized by the crustose thallus, a usually distinct prothallus, black lecideine apothecia, *Rhizocarpon*-type and (1–)8-spored asci, and ellipsoid halonate hyaline to brown or olive-green ascospores that can be transversely septate or submuriform to muriform; *Rhizocarpon* species usually contain secondary compounds detectable by TLC (Fletcher & al. 2009; McCarthy & Elix 2014). *Rhizocarpon* includes about 226 species worldwide (Lücking & al. 2016, Davydov & Yakovchenko 2017), of which 44 species have been reported from China (Wei 1991; Abbas & Wu 1998; Aptroot & Sparrius 2003; Li & al. 2013; Zhao & al. 2013; Mahire & al. 2015; Wang 2015a,b,c, 2016). The genus is most diverse and abundant

* Ling Hu & Xiao Zhang contributed equally to this research.

on siliceous rocks, with some species on calcareous rocks and a minority parasitic on other lichens in montane habitats and at temperate to higher latitudes (Ihlen 2004, McCarthy & Elix 2014).

This paper contributes further to our understanding of *Rhizocarpon* in China in the context of the Lichen Flora of China project. We have identified four species new to the country: *R. cinereonigrum*, *R. distinctum*, *R. sublavatum* and *R. subpostumum*.

Materials & methods

The specimens documented here are deposited in the Lichen Section of Botanical Herbarium, Shandong Normal University, Jinan, China (SDNU). Morphological and anatomical characters were examined under a COIC XTL7045B2 stereomicroscope and Olympus CX41 polarizing microscope. The thallus and medulla were tested with K (a 10% aqueous solution of potassium hydroxide), C (a solution of aqueous sodium hypochlorite), and I (Lugol's iodine) for identification. Lichen substances were identified using standardized thin-layer chromatography techniques (TLC) with solvent system C (Orange & al. 2010). The lichens were photographed with an Olympus SZX16 and a BX61 microscope with a DP72 lens.

Taxonomy

Rhizocarpon cinereonigrum Vain.,

Acta Soc. Fauna Flora Fenn. 53(1): 279 (1922)

FIG. 1

THALLUS crustose, with plane, contiguous areolae, dark greyish to dark brown, esorediate. PROTHALLUS absent. MEDULLA white, I–. APOTHECIA black, round to rounded-angular, 0.3–0.5 mm diam., immersed to sessile, lecideine with a well-developed exciple, persistent, concave to flat. PROPER EXCIPLE dark brown, with crystals partly dissolving in K; epihymenium dark brown, K+ violet-red; hymenium hyaline; paraphyses branched and anastomosing; hypothecium dark brown, K–. ASCI 8-spored. ASCOSPORES soon brownish green, 1-septate, with a halo, constricted at the septum when overmature, $28\text{--}38 \times 12\text{--}18 \mu\text{m}$. CONIDIOMATA not seen.

CHEMISTRY—Medulla K+ yellow, C–. Stictic acid detected by TLC.

SPECIMENS EXAMINED: CHINA. SICHUAN, Kangding Co., Mt Zheduo, alt. 4200 m, on siliceous rock, 15 Oct. 2015, Weicheng Wang 20150702 (SDNU); Luding Co., Moxi Town, Yajiagen, alt. 4000 m, on siliceous rock, 13 Oct. 2015, Weicheng Wang 20150665 (SDNU). YUNNAN, Shangri-La, Mt Hong, alt. 4462 m, on siliceous rock, 18 Aug. 2018, Chunxiao Wang 20180633 (SDNU).

DISTRIBUTION—Finland, Novaya Zemlya, and Greenland (Lynge 1932; Thomson 1997). New to China.




FIG. 1. *Rhizocarpon cinereonigrum* (SDNU – Wang 20150665). A: Thallus; B: Apothecia; C: Apothecium section; D: Crystals in exciple; E: K reaction; F: Ascospores.

COMMENTS—Our specimens are similar to those described previously in the literature, except that the protologue description mentions a prothallus and convex apothecia (Thomson 1967, 1997). *Rhizocarpon cinereonigrum* is characterized by the dark greyish to dark brown thallus, epihymenium K+ violet-red, large 1-septate ascospores, and containing stictic acid. This species morphologically resembles *R. badioatrum* (Flörke ex Spreng.) Th. Fr. and

R. jemtlandicum Malme; however, *R. badioatrum* differs by containing $\pm$ stictic, $\pm$ norstictic; and diffractic acids, while *R. jemtlandicum* differs in its K–, olive to green-black epithecium (Timdal & Holtan-Hartwig 1988, Fletcher & al. 2009, Wirth & al. 2013).




FIG. 2. *Rhizocarpon distinctum* (SDNU – Wang 20151607). A: Thallus; B: Apothecia; C: Apothecium section; D: Crystals in epihymenium; E: Ascus; F: Ascospores.

Rhizocarpon distinctum Th. Fr., Lichenogr. Scand. 2: 625 (1874) FIG. 2
THALLUS areolate; areoles small, dark grey, contiguous or more rarely scattered, rounded to angular, convex. PROTHALLUS black. MEDULLA white,

I+ blue. APOTHECIA lecideine, black, 0.3–0.5 mm diam., sessile, concave to flat; exciple thin, dark brown, persistent; epihymenium brown to olive-brown, K–, with crystals partly dissolving in K; hymenium hyaline; hypothecium dark brown. ASCI 8-spored. ASCOSPORES soon brown, 3-septate to submuriform with 4–6 cells, with a halo, $20\text{--}27.5 \times 7.5\text{--}12.5\ \mu\text{m}$. CONIDIOMATA not seen.

CHEMISTRY—Medulla K+ yellow, C–. Stictic acid detected by TLC.

SPECIMEN EXAMINED: CHINA. SICHUAN, Mt. Zheduo, alt. 4000 m, on siliceous rocks, 15 Oct. 2015, Weicheng Wang 20151607 (SDNU).

DISTRIBUTION—*Rhizocarpon distinctum* has been reported from Europe, British Isles, North America (eastern Canada), Antarctica, Novaya Zemlya, Australia, and New Zealand (Lynge 1932, Thomson 1997, Øvstedal & Smith 2001, Ihlen 2004, Fletcher & al. 2009, Galloway 2007, McCarthy & Elix 2014). New to China.

COMMENTS—Our specimens match previously published descriptions, except for the presence of an epihymenium containing crystals partly dissolving in K (Ihlen 2004, Fletcher & al. 2009). *Rhizocarpon distinctum* is characterized by the amyloid medulla, 3-septate to submuriform ascospores, and the occurrence of stictic acid (Timdal & Holtan-Hartwig 1988, Thomson 1997, Øvstedal & Smith 2001, Ihlen 2004, Fletcher & al. 2009, McCarthy & Elix 2014). The small areoles and apothecia of *R. distinctum* are suggestive of *R. reductum* Th. Fr., *R. sublavatum* and *R. timdalii* Ihlen & Fryday, but *R. distinctum* can be distinguished from those species in becoming brown and in usually having fewer than eight cells per ascospore (Ihlen 2004, Fletcher & al. 2009).

Rhizocarpon sublavatum Fryday, Lichenologist 32(3): 214 (2000)

FIG. 3

THALLUS areolate, greyish brown, continuous, flat. PROTHALLUS black. MEDULLA white, I–. APOTHECIA lecideine, black, round, 0.3–0.6 mm diam., sessile, concave; disc epruinose; exciple black, thick, usually persistent. PROPER EXCIPLE dark brown, without crystals; epihymenium bluish green, K+ brighter; hymenium hyaline; hypothecium dark brown. ASCI 8-spored. ASCOSPORES colourless, eumuriform, $25\text{--}35\text{--}(37.5) \times (10\text{--})12.5\text{--}15\text{--}(17.5)\ \mu\text{m}$. CONIDIOMATA not seen.

CHEMISTRY—Thallus K–, C–. Medulla K–, C–. No substances detected by TLC.

SPECIMEN EXAMINED: CHINA. SICHUAN, Luding Co., Moxi Town, Park. Hongshi, alt. 3060 m, on siliceous rock, 13 Oct. 2015, Weicheng Wang 20150660 (SDNU)




FIG. 3. *Rhizocarpon sublavatum* (SDNU – Wang 20150660). A: Thallus; B: Apothecia; C: Apothecium section; D: Ascospores.

DISTRIBUTION—Great Britain (Scotland), Scandinavia, Iceland, and Central and Eastern Europe (Fryday 2000; Ihlen 2004). New to China.

COMMENTS—Our specimen agrees with the type description, except that the protologue cites an effuse thallus and a flat to slightly convex disc (Fryday 2000). *Rhizocarpon sublavatum* is characterized by having smaller eumuriform spores than *R. lavatum* (Ach.) Hazsl., a bluish green epihymenium, and no lichen substances. The species shares a similar ascospore size and shape with *R. anaperum* (Vain.) Vain., which differs in a browner, more granular-areolate thallus, more highly branched paraphysoids, and a different apothecial pigmentation with only brown and grey pigments (Fryday 2000).

Rhizocarpon subpostumum (Nyl.) Arnold,
Verh. Zool.-Bot. Ges. Wien 27: 554 (1877)

FIG. 4

THALLUS areolate; areoles flat to slightly convex, polygonal to rounded, brown, epruinose. **PROTHALLUS** indistinct. **MEDULLA** white, I–. **APOTHECIA**




FIG. 4. *Rhizocarpon subpostumum* (SDNU – Du 20080268). A: Thallus and apothecia; B: Apothecium section; C: K reaction; D: Ascospores.

black, rounded to angular, 0.3–0.6(–0.8) mm diam., usually immersed, lecideine with a thin margin, flat to convex. PROPER EXCIPIE dark brown, K+ purple; epihymenium dark green-black, K–; hymenium vertically streaked red-brown and K+ purple; hypothecium dark brown, K–. ASCI 8-spored. ASCOSPORES colorless, pale brown green when over-mature, submuriform, with a halo, $15\text{--}22.5\text{--}(27.5) \times 7.5\text{--}12.5\text{--}(14) \mu\text{m}$. CONIDIOMATA not seen.

CHEMISTRY—Thallus K–, C–; medulla K–, C–. No substances detected by TLC.

SPECIMENS EXAMINED: CHINA. SICHUAN, Litang, Mt Kazila, alt. 4700 m, on siliceous rock, 7 Nov. 2008, Yuanda Du 20080111, 20180268 (SDNU).

DISTRIBUTION—*Rhizocarpon subpostumum* has previously been reported from Norway, Sweden, Finland, Iceland, Great Britain (Scottish Highlands), North America, and New Zealand (Ihlen 2004, Galloway 2007, Fletcher & al. 2009). New to China.

COMMENTS—Our specimens conform to previously published descriptions. *Rhizocarpon subpostumum* is characterized by small apothecia, small submuriform ascospores, and the K+ purple exciple. It shares small apothecia and ascospores with *R. postumum* (Nyl.) Arnold, which is distinguished by its exciple not reacting K+ purple (Fryday 2000, Ihlen 2004, Fletcher & al. 2009).

Acknowledgements

The authors thank Dr. Patrick McCarthy (Canberra, A.C.T., Australia) and Dr. Shou-Yu Guo (State Key Laboratory of Mycology, Institute of Microbiology, Chinese Academy of Sciences, Beijing, China) for guidance in specimen identification and pre-submission review. We also thank the others in our laboratory for assistance during this study. This investigation was supported by the Emergency management project of National Natural Science Foundation of China (31750001), the National Natural Science Foundation of China (31900010) and the Scientific Research Foundation of Graduate School of Shandong Normal University (SCX201945).

Literature cited

- Abbas A, Wu JN. 1998. Lichens of Xinjiang. Sci-Tech & Hygiene Publishing House of Xinjiang (K), Urumqi. 178 p.
- Aptroot A, Sparrius LB. 2003. New microlichens from Taiwan. Fungal Diversity 14: 1–50.
- Davydov EA, Yakovchenko LS. 2017. *Rhizocarpon smaragdulum*, a new monosporic yellow-thalline species and some additional species of the genus *Rhizocarpon* from the Altai Mountains (Siberia). Lichenologist 49(5): 457–466.
<https://doi.org/10.1017/S0024282917000469>
- Fletcher A, Gilbert OL, Clayden S, Fryday AM. 2009. *Rhizocarpon* Ramond ex DC. (1805). 792–808, in: CW Smith & al. (eds). Lichens of Great Britain and Ireland. British Lichen Society, London.
- Fryday A. 2000. On *Rhizocarpon obscuratum* (Ach.) Massal., with notes on some related species in the British Isles. Lichenologist 32(3): 207–224. <https://doi.org/10.1006/lich.2000.0269>
- Galloway DJ. 2007. *Rhizocarpon* Ramond ex DC. 1529–1555, in: Flora of New Zealand lichens, revised 2nd ed., vol. 2: *Pannaria*–**Zwackhiomyces*. Manaaki Whenua Press, Lincoln, New Zealand.
- Ihlen PG. 2004. Taxonomy of the non-yellow species of *Rhizocarpon* (*Rhizocarpaceae*, lichenized *Ascomycota*) in the Nordic countries, with hyaline and muriform ascospores. Mycological Research 108: 533–570. <https://doi.org/10.1017/S0953756204009803>
- Li X, Li C, Wang HY. 2013. Two species of *Rhizocarpon* new to China. Modern Agricultural Science and Technology 6: 146–147.
- Lücking R, Hodkinson BP, Leavitt SD. 2016. The 2016 classification of lichenized fungi in the *Ascomycota* and *Basidiomycota*: Approaching one thousand genera. Bryologist 119(4): 361–416. <https://doi.org/10.1639/0007-2745-119.4.361>
- Lyngbe B. 1932. A revision of the genus *Rhizocarpon* (Ram.) T. Fr. in Greenland. Skrifer om Svalbard og Ishavet 47: 1–30.

- McCarthy PM, Elix JA. 2014. The lichen genus *Rhizocarpon* in mainland Australia. *Telopea* 16: 195–211. <https://doi.org/10.7751/telopea20148124>
- Øvstedal DO, Smith RIL. 2001. Lichens of Antarctica and South Georgia. Cambridge University Press, Cambridge.
- Timdal E, Holtan-Hartwig J. 1988. A preliminary key to *Rhizocarpon* in Scandinavia. *Graphis Scripta* 2: 41–54.
- Thomson JW. 1967. Notes on *Rhizocarpon* in the Arctic. *Nova Hedwigia* 14: 421–481.
- Thomson JW. 1997. American Arctic lichens 2. Microlichens. Madison: University of Wisconsin Press. 675 p.
- Wang WC, Zhao ZT. 2015a. Four new records of *Rhizocarpon* from China. *Mycotaxon* 130: 739–747. <https://doi.org/10.5248/130.739>
- Wang WC, Zhao ZT. 2015b. Four *Rhizocarpon* species new to China. *Mycotaxon* 130: 883–891. <https://doi.org/10.5248/130.883>
- Wang WC, Zhao ZT. 2015c. Three new records of *Rhizocarpon* from China. *Acta Botanica Boreali-Occidentalia Sinica* 35(8): 1694–1696.
- Wang WC, Ren ZJ, Zhang LL. 2016. New records of *Rhizocarpon* from Hengduan Mountains, China. *Mycotaxon* 131: 589–596. <https://doi.org/10.5248/131.589>
- Wei JC. 1991. An enumeration of lichens in China. International Academic Publishers, Beijing.
- Wirth V, Hauck M, Schultz M. 2013. Die Flechten Deutschlands, Band 2. E. Ulmer, Stuttgart.
- Zhao ZT, Li C, Zhao X, Zhang LL. 2013. New records of *Rhizocarpon* from China. *Mycotaxon* 125: 217–226. <https://doi.org/10.5248/125.217>